

Biochimie de la dépression et des antidépresseurs

M. STROLIN BENEDETTI

Résumé. Les voies de synthèse et de dégradation des principales monoamines (dopamine, noradrénaline, sérotonine, phényléthylamine) impliquées dans la pathologie des dépressions sont exposées rapidement. Les variations de l'activité des enzymes responsables des taux sérique, urinaire et cérébrospinal des monoamines et de leurs principaux métabolites sont discutées en relation avec les différents états dépressifs. La modification de la sensibilité des récepteurs neuroaux dans la dépression est ensuite abordée.

Les effets biochimiques spécifiques des inhibiteurs de la monoamine oxydase (IMAO) et des inhibiteurs de la recapture des monoamines sont exposés. Les IMAO sont divisés en inhibiteurs mixtes (A + B) et irréversibles, inhibiteurs spécifiques et irréversibles, inhibiteurs spécifiques et réversibles et leur mécanisme d'action avec l'enzyme est également présenté. Les inhibiteurs de la recapture des monoamines sont séparés selon leur spécificité d'action vis-à-vis des monoamines. Le mode d'action de certains antidépresseurs considérés comme atypiques est également abordé. Enfin, l'effet d'un traitement chronique par les antidépresseurs sur la sensibilité des récepteurs noradrénergiques, sérotoninergiques et dopaminergiques pré- et post-synaptiques est discuté.

*Biochemistry of depression and of antidepressant action.*M. STROLIN BENEDETTI. *L'Encéphale*, 1982, VIII, 545-585.

Summary. The pathways for the synthesis and degradation of the principal monoamines (dopamine, noradrenaline, serotonin, and phenylethylamine) concerned in the pathology of depression are briefly presented. Differences in the activities of the enzymes which control the amounts of monoamines or their principal metabolites in serum, urine or cerebrospinal fluid are discussed in relation to the different types of depression, as is also the role of changes in the sensitivity of neuronal receptors. The specific biochemical effects of monoamine oxidase inhibitors (MAOI's) and of the inhibitors of monoamine re-uptake are considered. MAOI's are classified as irreversible mixed inhibitors (A + B), as irreversible specific inhibitors and as reversible specific inhibitors; their mechanism of interaction with the enzyme is presented. Inhibitors of monoamine re-uptake are classified according to their specificity for different monoamines. The mechanism of action of some antidepressants which have an atypical mechanism of action is outlined. Finally, the effect of chronic treatment with antidepressants on the sensitivities of the noradrenergic, serotonergic and dopaminergic pre- and post-synaptic receptors is discussed.


Travail reçu le 7 juin 1982 et accepté le 21 juin 1982.

Tirés à part: M. STROLIN BENEDETTI, Centre de Recherche Delalande, 10, rue des Carrières, F 92500 Rueil-Malmaison.

Visez l'efficacité.

Composition - Forme Présentation. Vivalan 100 : boîte de vingt comprimés sous blister, dosés à 100 mg de viloxazine base.

Indications pharmacologiques. Antidépresseur. Indications : états dépressifs quel que soit le type : psychotique, névrotique ou réactionnel. Dépression psychotique et mélancolique, dépression de la psychose manico-dépressive, mélancolie d'invololution, psychoses dépressives réactionnelles, dépression névrotique ou réactionnelle, syndrome subjectif post-traumatique à manifestation dépressive, dépression céphalalgiques. Effets indésirables : Quelques survenues de nausées, diarrées, céphalées, troubles du sommeil, accélération modérée de la vitesse de sédimentation, très rarement modifications de l'E.C.G. et de l'E.E.G. Précautions : L'absorption d'alcool est déconseillée pendant la durée du traitement, notamment chez les éthyliques, la tolérance du Vivalan se trouvant alors diminuée. Interactions médicamenteuses : Une diminution de la posologie de la phénytoïne peut être nécessaire en cas de co-prescription. Propriétés : États dépressifs de faible gravité : 300 mg/jour. États dépressifs plus sévères : 800 à 600 mg/jour. La prescription doit se faire en deux ou trois prises quotidiennes, la dernière avant 17 h. En traitement d'entretien, il est possible d'utiliser le Vivalan 50. Elle peut être accompagnée d'une co-prescription d'anxiolytiques et éventuellement d'un hypotenseur. Vivalan 100. Thibaut A.M.M. 319700A. Prix : 36,60 F + S.T.P.A. Remb. S.S. à 70 % et C.G. Cati. moyen journalier de traitement : 5,56 F pour 800 mg/jour. I.C.I. Pharma 6, rue Blanche 95800 Enghien. Tél. 412.90.60.


VIVALAN 100
viloxazine

Antidépresseur majeur d'utilisation quotidienne

BEST AVAILABLE COPY

HARRISON'S 15TH EDITION

PRINCIPLES OF INTERNAL MEDICINE

EDITORS

EUGENE BRAUNWALD, MD,
MD(Hon), ScD(Hon)

Distinguished Hersey Professor of Medicine,
Faculty Dean for Academic Programs at Brigham and
Women's Hospital and Massachusetts General Hospital,
Harvard Medical School; Vice-President for Academic
Programs, Partners HealthCare Systems, Boston

ANTHONY S. FAUCI, MD,
ScD(Hon)

Chief, Laboratory of Immunoregulation; Director,
National Institute of Allergy and Infectious Diseases,
National Institutes of Health, Bethesda

DENNIS L. KASPER, MD,
MA(Hon)

William Ellery Channing Professor of Medicine,
Professor of Microbiology and Molecular Genetics,
Executive Dean for Academic Programs, Harvard
Medical School; Director, Channing Laboratory,
Department of Medicine, Brigham and Women's
Hospital, Boston

STEPHEN L. HAUSER, MD

Betty Anker Fife Professor and Chairman,
Department of Neurology,
University of California San Francisco,
San Francisco

DAN L. LONGO, MD

Scientific Director, National Institute on Aging,
National Institutes of Health,
Bethesda and Baltimore

J. LARRY JAMESON, MD, PhD

Irving S. Cutter Professor and Chairman,
Department of Medicine,
Northwestern University Medical School;
Physician-in-Chief, Northwestern
Memorial Hospital, Chicago

McGraw-Hill

MEDICAL PUBLISHING DIVISION

New York San Francisco Washington, DC Auckland Bogotá Caracas Lisbon London
Madrid Mexico City Milan Montreal New Delhi San Juan Singapore Sydney Tokyo Toronto



Note: Dr. Fauci and Dr. Longo's works as editors and authors were performed outside the scope of their employment as U.S. government employees. These works represent their personal and professional views and not necessarily those of the U.S. government.

**Harrison's
PRINCIPLES OF INTERNAL MEDICINE
Fifteenth Edition**

Copyright © 2001, 1998, 1994, 1991, 1987, 1983, 1980, 1977, 1974, 1970, 1966, 1962, 1958 by The McGraw-Hill Companies, Inc. All rights reserved. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a data base or retrieval system, without the prior written permission of the publisher.

1234567890 DOWDOW 0987654321

ISBN 0-07-007272-8 (Combo)

0-07-007273-6 (Vol. 1)

0-07-007274-4 (Vol. 2)

0-07-913686-9 (Set)

FOREIGN LANGUAGE EDITIONS

Arabic (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)

Chinese (Twelfth Edition)—McGraw-Hill Book Company—Singapore © 1994

Croatian (Thirteenth Edition)—Placebo, Split, Croatia

French (Fourteenth Edition)—McGraw-Hill Publishing Co., Maidenhead, UK © 1999

German (Fourteenth Edition)—McGraw-Hill Publishing Co., Maidenhead, UK © 1999

Greek (Fourteenth Edition)—Parissianos, Athens, Greece © 2000

Italian (Fourteenth Edition)—McGraw-Hill Libri Italia srl, Milan © 1999

Japanese (Eleventh Edition)—Hirokawa © 1991

Polish (Fourteenth Edition)—Czelej Publishing Company, Lubin, Poland (est. 2000)

Portuguese (Fourteenth Edition)—McGraw-Hill Interamericana do Brasil Ltda © 1998

Romania (Fourteenth Edition)—Teora Publishers, Bucharest, Romania (est. 2000)

Spanish (Fourteenth Edition)—McGraw-Hill Interamericana de Espana, Madrid © 1998

Turkish (Thirteenth Edition)—McGraw-Hill Libri Italia srl (est. 1996)

This book was set in Times Roman by Progressive Information Technologies. The editors were Martin Wonsiewicz and Mariapaz Ramos Englis. The production director was Robert Laffler. The index was prepared by Irving C. Tullar. The text and cover designer was Marsha Cohen/Parallelogram Graphics.

R. R. Donnelley and Sons, Inc. was the printer and binder.

Library of Congress Cataloging-in-Publication Data

Harrison's principles of internal medicine—15th ed./editors, Eugene Braunwald . . . [et al.]. p. cm.

Includes bibliographical references and index.

ISBN 0-07-913686-9 (set)—ISBN 0-07-007273-6 (v. 1)—ISBN 0-07-007274-4 (v. 2)

1. Internal medicine. I. Braunwald, Eugene, date

RC46.H333 2001

616—dc21

00-063809

BEST AVAILABLE COPY

Table 385-3 Antidepressants

Name	Usual Daily Dose, mg	Side Effects	Comments
SSRIs			
Fluoxetine (Prozac)	10-80	Headache; nausea and other GI effects; jitteriness; insomnia; sexual dysfunction; can affect plasma levels of other meds (except sertraline); akathisia rare	Once daily dosing, usually in A.M.; fluoxetine has very long half-life; must not be combined with MAOIs
Sertraline (Zoloft)	50-200		
Paroxetine (Paxil)	20-60		
Fluvoxamine (Luvox)	100-300		
Citalopram (Celexa)	20-60		
TCAs			
Amitriptyline (Elavil)	150-300	Anticholinergic (dry mouth, tachycardia, constipation, urinary retention, blurred vision); sweating; tremor; postural hypotension; cardiac conduction delay; sedation; weight gain	Once daily dosing, usually qhs; blood levels of most TCAs available; can be lethal in O.D. (lethal dose = 2 g); nortriptyline best tolerated, especially by elderly
Nortriptyline (Pamelor)	50-200		
Imipramine (Tofranil)	150-300		
Desipramine (Norpramin)	150-300		
Doxepin (Sinequan)	150-300		
Clomipramine (Anafranil)	150-300		
Mixed norepinephrine/serotonin reuptake inhibitors			
Venlafaxine (Effexor)	75-375	Nausea; dizziness; dry mouth; headaches; increased blood pressure; anxiety and insomnia	Bid-tid dosing; lower potential for drug-drug interactions than SSRIs; contraindicated with MAOIs. Once daily dosing
Mirtazapine (Remeron)	15-45	Somnolence; weight gain; neutropenia rare	
Mixed-action drugs			
Bupropion (Wellbutrin)	250-450	Jitteriness; flushing; seizures in at-risk patients; anorexia; tachycardia; psychosis	Tid dosing, but sustained release also available; fewer sexual side effects than SSRIs or TCAs; may be useful for adult ADD
Trazodone (Desyrel)	200-600	Sedation; dry mouth; ventricular irritability; postural hypotension; priapism rare	Useful in low doses for sleep because of sedating effects with no anticholinergic side effects
Nefazodone (Serzone)	300-600	Sedation; headache; dry mouth; nausea; constipation	Once daily dosing; no effect on REM sleep unlike other antidepressants
MAOIs			
Phenelzine (Nardil)	45-90	Insomnia; hypotension; anorgasmia; weight gain; hypertensive crisis; tyramine cheese reaction; lethal reactions with SSRIs; serious reactions with narcotics	May be more effective in patients with atypical features or treatment-refractory depressions
Tranylcypromine (Parnate)	20-50		
Isocarboxazid (Marplan)	20-60		

NOTE: ADD, attention deficit disorder; MAOI, monoamine oxidase inhibitor; REM, rapid eye movement; SSRI, selective serotonin reuptake inhibitor; TCA, tricyclic antidepressant.

autonomic responsivity, and social learning. Panic disorder shows familial aggregation, although concordance in monozygotic twins is only 30%. Acute panic attacks appear to be associated with increased noradrenergic discharge in the locus coeruleus. Intravenous infusion of sodium lactate evokes an attack in two-thirds of panic disorder patients, as do the α_2 -adrenergic antagonist yohimbine and carbon dioxide inhalation. It is hypothesized that each of these stimuli activates a neural circuit involving noradrenergic neurons in the locus coeruleus and serotonergic neurons in the dorsal raphe. Agents that block serotonin reuptake are therapeutic in preventing attacks. It is theorized that panic-disorder patients have a heightened sensitivity to somatic symptoms, which triggers increasing arousal, setting off the "panic attack" mechanism. Accordingly, successful therapeutic intervention involves altering the patient's cognitive interpretation of anxiety-producing experiences as well as preventing the attack itself.

TREATMENT Achievable goals of treatment are to decrease the frequency of panic attacks and to reduce their intensity. The cornerstone of drug therapy is antidepressant medications (Tables

385-3, 385-4, and 385-5). T antidepressant (TCA) agents imi clomipramine can benefit 75 panic disorder patients. Low dose 25 mg/d) are given initially to a creased anxiety associated with monoamine levels in the initi treatment. Selective serotonin reu tors (SSRIs) are equally effective have the adverse effects of T should be started at one-third to their usual antidepressant dose (mg fluoxetine, 25 to 50 mg sertr paroxetine). Monoamine oxidase (MAOIs) are at least as effective and may specifically benefit patients have comorbid features of atypical (i.e., hypersomnia and weight gain orthostatic hypotension, and to maintain a low-tyramine diet (i cheese and wine) have limited the ever. Antidepressants typically weeks to become effective, and need to be adjusted according to sponse.

Because of anticipatory anxiety need for immediate relief of panic benzodiazepines are useful early of treatment and sporadically then 385-6). For example, alprazolam 0.5 mg qid and increasing to 4 mg doses, is effective, but patients must be monitored closely, as some develop dependence begin to escalate the dose of this Clonazepam, at a final maintenance to 4 mg/d, is also helpful; its long permits twice-daily scheduling, appear less likely to develop dependence this agent.

Early psychotherapeutic intervention psychoeducation aimed at symptom hances the effectiveness of drug treatment. Patients can be taught breathing techniques be educated about physiologic changes occur with panic, and can learn to control themselves voluntarily to precipitate Homework assignments and medication compliance are important components of full treatment. Once patients have

satisfactory response, drug treatment should be maintained for years to prevent relapse.

GENERALIZED ANXIETY DISORDER **Characteristics** Patients with generalized anxiety disorder persistent, excessive, and/or unrealistic worry associated signs and symptoms, which commonly include muscle paired concentration, autonomic arousal, feeling "on edge and insomnia (Table 385-7). Onset is usually before history of childhood fears and social inhibition may be incidence of GAD is increased in first-degree relatives of the diagnosis; family studies also indicate that GAD and segregate independently. Over 80% of patients with GAD from major depression, dysthymia, or social phobia. Substance abuse is common in these patients, particularly sedative/hypnotic abuse. Patients with GAD readily admit excessively over minor matters, with life-disrupting effects panic disorder, complaints of symptoms such as shortness palpitations, and tachycardia are relatively rare.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ ~~BLACK BORDERS~~

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ ~~FADED TEXT OR DRAWING~~

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.